

JOURNAL OF ALGEBRA **151**, 1–11 (1992)

On the Antipode of a Quasitriangular Hopf Algebra

DAVID E. RADFORD

*Department of Mathematics, Statistics, and Computer Science,
University of Illinois at Chicago, Chicago, Illinois 60680*

Communicated by Nathan Jacobson

Received October 1, 1989

Let (A, R) be a quasitriangular Hopf algebra with antipode s in the category of vector spaces over a field k . In this paper we show that s^2 is inner, which means s is bijective, and when A is finite-dimensional there is a grouplike element $h \in A$ such that $s^4(a) = hah^{-1}$ for all $a \in A$.

Suppose that A is any Hopf algebra with antipode s over the field k . Proposition 1 of Section 1 gives a sufficient condition for s^2 to be inner which we apply to quasitriangular Hopf algebras in Section 2. Its statement and proof are based on calculations found in [2, pp. 66–67].

Generally s^2 need not be inner. Suppose that A is finite-dimensional. If s^2 were inner, then all ideals of A would be invariant under s^2 , and hence all subcoalgebras of the dual Hopf algebra A^* would be invariant under S^2 , where $S = s^*$ is the antipode of A^* . There are examples [5] over algebraically closed fields in which the latter is not the case.

Section 2 begins with a discussion of properties of quasitriangular Hopf algebras (A, R) over the field k used in this paper. Write $R = \sum R^{(1)} \otimes R^{(2)} \in A \otimes A$ and set $u = \sum s(R^{(2)}) R^{(1)}$. The main result of Section 2 is that u is invertible and $s^2(a) = uau^{-1}$ for all $a \in A$. This result was established in [2, Exercise 7.3.6] under the hypothesis that s is bijective.

If A is any finite-dimensional Hopf algebra with antipode s over the field k , there are distinguished grouplike elements $g \in A$ and $\alpha \in A^*$ which relate left and right integrals and which together describe s^4 . In Section 3 we consider s^4 when (A, R) is a finite-dimensional quasitriangular Hopf algebra. Let $h = vu$, where $u = \sum s(R^{(2)}) R^{(1)}$ is defined as above and $v = s(u)^{-1}$. By relating h to g and α , we are able to show that h is a grouplike element and that $s^4(a) = hah^{-1}$ for all $a \in A$. When A is unimodular we prove that $h = g$. Thus the product vu does not depend on R in the unimodular case.

We assume that the reader has some familiarity with the elementary aspects of the theory of Hopf algebras. A recommended reference is [6].

1. A SUFFICIENT CONDITION FOR s^2 TO BE INNER

Suppose that A is a Hopf algebra with antipode s over the field k . Let $\Delta^{cop}(a) = \sum a_{(2)} \otimes a_{(1)}$ for $a \in A$ denote the coproduct on A derived by “twisting” comultiplication, and let A^{cop} denote the resulting bialgebra. Define a right module action of A on itself by

$$b \triangleleft a = \sum s(a_{(2)}) b a_{(1)} \quad \text{for } a \in A.$$

Observe that

$$\begin{aligned} s^2(a) b &= \sum s^2(a_{(1)}) b (\varepsilon(a_{(2)}) 1) \\ &= \sum s^2(a_{(1)}) b s(a_{(2)}) a_{(3)} \\ &= \sum s((s(a_{(1)}))_{(2)}) b (s(a_{(1)}))_{(1)} a_{(2)} \\ &= \sum (b \triangleleft s(a_{(1)})) a_{(2)} \end{aligned}$$

for all $a, b \in A$ since s is a coalgebra antihomomorphism [6, Proposition 4.0.1]. Thus left and right multiplication in A are related by

$$s^2(a) b = \sum (b \triangleleft s(a_{(1)})) a_{(2)} \quad \text{for all } a, b \in A. \quad (1)$$

PROPOSITION 1. *Suppose that A is a Hopf algebra with antipode s over the field k . Suppose further that $R = \sum R^{(1)} \otimes R^{(2)} \in s(A) \otimes A$ has an inverse in $A \otimes A$, and that $\Delta^{cop}(a) = R(\Delta(a)) R^{-1}$ for all $a \in A$. Then $u = \sum s(R^{(2)}) R^{(1)}$ is invertible and $s^2(a) = uau^{-1}$ for all $a \in A$. In particular s is bijective.*

Proof. Let $a \in A$. We first consider implications of the hypothesis $\Delta^{cop}(a) = R(\Delta(a)) R^{-1}$, which can be formulated $(\Delta^{cop}(a)) R = R(\Delta(a))$.

Suppose that $X = \sum X^{(1)} \otimes X^{(2)} \in A \otimes A$ satisfies $(\Delta^{cop}(a)) X = X(\Delta(a))$ for all $a \in A$, which is the same as

$$\sum a_{(2)} X^{(1)} \otimes a_{(1)} X^{(2)} = \sum X^{(1)} a_{(1)} \otimes X^{(2)} a_{(2)} \quad \text{for all } a \in A. \quad (2)$$

Let $x = \sum s(X^{(2)}) X^{(1)}$. Noting that s is an algebra antihomomorphism [6, Proposition 4.0.1], we apply $I \otimes s$ to both sides of Eq. (2), exchange tensors, and then multiply to obtain

$$x \triangleleft a = \varepsilon(a) x \quad \text{for all } a \in A. \quad (3)$$

By Eqs. (1) and (3)

$$s^2(a) x = xa \quad \text{for all } a \in A. \quad (4)$$

Write $R^{-1} = \sum U^{(1)} \otimes U^{(2)}$. Since $R^{-1}(\Delta^{cop}(a)) = (\Delta(a)) R^{-1}$ for all $a \in A$, we have that $\sum U^{(1)} a_{(2)} \otimes U^{(2)} a_{(1)} = \sum a_{(1)} U^{(1)} \otimes a_{(2)} U^{(2)}$, and hence $\sum a_{(2)} U^{(2)} \otimes a_{(1)} U^{(1)} = \sum U^{(2)} a_{(1)} \otimes U^{(1)} a_{(2)}$ for all $a \in A$. Set $v = \sum s(U^{(1)}) U^{(2)}$. Then by Eqs. (2) and (4) we conclude that

$$s^2(a) u = ua \quad \text{and} \quad s^2(a) v = va \quad \text{for all } a \in A. \quad (5)$$

Now $\sum U^{(1)} R^{(1)} \otimes U^{(2)} R^{(2)} = 1 \otimes 1$. Applying $s \otimes I$ to this equation and multiplying, we obtain $\sum s(R^{(1)}) s(U^{(1)}) U^{(2)} R^{(2)} = 1$, or $\sum s(R^{(1)}) v R^{(2)} = 1$. Therefore by Eq. (5) we have that $\sum s(R^{(1)}) s^2(R^{(2)}) v = 1$, or equivalently

$$s(u) v = 1. \quad (6)$$

By Eq. (6) it follows that v has a left inverse, and hence s is injective by Eq. (5).

Now we use the hypothesis $R \in s(A) \otimes A$ to show that v has a right inverse. Under this hypothesis $u = \sum s(R^{(2)}) R^{(1)} \in s(A)$, so $u = s(w)$ for some $w \in A$. Thus by Eqs. (6) and (5) it follows that $1 = s(u) v = s^2(w) v = vw$. Therefore v is invertible, and hence $s^2(a) = vav^{-1}$ for all $a \in A$ by Eq. (5). In particular s is bijective. This implies that u is invertible also, and hence $s^2(a) = uau^{-1}$ for all $a \in A$ by Eq. (5) again. The proof of the proposition is complete.

An easy corollary to the proof is that when A is finite-dimensional, and $R, U \in A \otimes A$ satisfy $\Delta^{cop}(a) = R(\Delta(a)) U$ for all $a \in A$, then s^2 is inner. The last proposition of Section 3 is an improvement of this observation.

2. QUASITRIANGULAR HOPF ALGEBRAS

In this section we show that the hypothesis of Proposition 1 is satisfied for a quasitriangular Hopf algebra (A, R) with antipode s in the category of vector spaces over the field k , and thus it will follow that s^2 is inner. We begin by giving a definition of quasitriangular Hopf algebras which is equivalent to the one of [1, p. 811] in our context.

DEFINITION. A *quasitriangular Hopf algebra* in the category of vector spaces over the field k is a pair (A, R) , where A is a Hopf algebra over k , and $R = \sum R^{(1)} \otimes R^{(2)} \in A \otimes A$ satisfies

$$(QT.1) \quad \sum \Delta R^{(1)} \otimes R^{(2)} = \sum R^{(1)} \otimes r^{(1)} \otimes R^{(2)} r^{(2)}, \text{ where } r = R,$$

$$(QT.2) \quad \sum \varepsilon(R^{(1)}) R^{(2)} = 1,$$

$$(QT.3) \quad \sum R^{(1)} \otimes \Delta R^{(2)} = \sum R^{(1)} r^{(1)} \otimes r^{(2)} \otimes R^{(2)}, \text{ where } r = R,$$

$$(QT.4) \quad \sum R^{(1)} \varepsilon(R^{(2)}) = 1, \text{ and}$$

$$(QT.5) \quad (\Delta^{cop}(a)) R = R(\Delta(a)) \text{ for all } a \in A.$$

Let A be a Hopf algebra over the field k , and let $F: A \otimes A \rightarrow \text{Hom}_k(A^*, A)$ be the (injective) linear map defined by $F(a \otimes b)(p) = p(a)b$ for all $a, b \in A$ and $p \in A^*$. Now suppose that $R = \sum R^{(1)} \otimes R^{(2)} \in A \otimes A$ and set $f = F(R)$.

It is easy to see that R satisfies (QT.1) and (QT.2) if and only if f is an algebra map. If f is an algebra map then R is invertible, for:

LEMMA 1. *Suppose that A is a Hopf algebra with antipode s over the field k , and $R \in A \otimes A$ satisfies (QT.1) and (QT.2). Then R is invertible with inverse $R^{-1} = \sum s(R^{(1)}) \otimes R^{(2)}$.*

Proof. First observe that (QT.2) implies that $1 \otimes 1 = \sum \varepsilon(R^{(1)}) 1 \otimes R^{(2)}$. By definition of the antipode $\sum s(a_{(1)}) a_{(2)} = \varepsilon(a) 1 = \sum a_{(1)} s(a_{(2)})$ for all $a \in A$. Now using (QT.1) it is easy to see that $1 \otimes 1 = \sum s(R^{(1)}) r^{(1)} \otimes R^{(2)} r^{(2)} = \sum R^{(1)} s(r^{(1)}) \otimes R^{(2)} r^{(2)}$. This concludes the proof of the lemma.

Suppose that R is invertible and that (QT.1) is satisfied. Then $\sum \varepsilon(R^{(1)}) R^{(2)}$ is an invertible idempotent, so (QT.2) is satisfied. Likewise if R is invertible and (QT.3) is satisfied, then (QT.4) is satisfied also. Thus the definition of quasitriangular Hopf algebras given above agrees with the one given in [1, p. 811] specialized to the category of vector spaces over k . From this point on we will refer to quasitriangular Hopf algebras in the category of vector spaces over k simply as quasitriangular Hopf algebras.

LEMMA 2. *Suppose that A is a Hopf algebra with antipode s over the field k , and $R \in A \otimes A$ satisfies (QT.3) and (QT.4). Then $\sum s(R^{(1)}) \otimes R^{(2)}$ is invertible with inverse $\sum s(R^{(1)}) \otimes s(R^{(2)})$. Thus $\sum R^{(1)} \otimes R^{(2)} = \sum s(R^{(1)}) \otimes s(R^{(2)})$ if (A, R) is quasitriangular.*

Proof. First observe that (QT.4) implies that $1 \otimes 1 = \sum R^{(1)} \otimes \varepsilon(R^{(2)}) 1$. Therefore (QT.3) and (QT.4) imply that $1 \otimes 1 = \sum R^{(1)} r^{(1)} \otimes s(r^{(2)}) R^{(2)} = \sum R^{(1)} r^{(1)} \otimes r^{(2)} s(R^{(2)})$. Since s is an algebra anti-homomorphism, applying $s \otimes I$ to these equations gives us that $\sum s(R^{(1)}) \otimes R^{(2)}$ is invertible with inverse $\sum s(R^{(1)}) \otimes s(R^{(2)})$. The last assertion now follows by Lemma 1. This completes the proof.

The two preceding lemmas are minor reformulations of material found in [2, p. 13].

Now suppose that A is finite-dimensional. Then it is easy to see that R satisfies (QT.3) and (QT.4) if and only if f is a coalgebra anti-homomorphism. In the finite-dimensional case F is an isomorphism of the algebra $A \otimes A$ and the convolution algebra $\text{Hom}_k(A^*, A^{cop})$. The Hopf algebra A^* has antipode $S = s^*$. Since A is finite-dimensional s is bijective

[6, Corollary 5.1.6]. Since s is bijective A^{cop} is a Hopf algebra with antipode s^{-1} .

Suppose further that (A, R) is quasitriangular. Then the conclusions of the lemmas have an easy explanation in terms of the convolution algebra $\text{Hom}_k(A^*, A^{cop})$. Since (A, R) is quasitriangular $f: A^* \rightarrow A^{cop}$ is a bialgebra homomorphism. Since f is an algebra homomorphism, f has an inverse in the convolution algebra which is $f \circ S$ by [6, Lemma 4.0.3.(1)]. Observe that $f \circ S$ corresponds to $\sum s(R^{(1)}) \otimes R^{(2)}$. Since f is a coalgebra homomorphism, f has an inverse in the convolution algebra which is $s^{-1} \circ f$ by [6, Lemma 4.0.3.(2)]. Therefore $f \circ S = s^{-1} \circ f$, or $f = s \circ f \circ S$. It is easy to see that $s \circ f \circ S$ corresponds to $\sum s(R^{(1)}) \otimes s(R^{(2)})$.

Now suppose that (A, R) is any quasitriangular Hopf algebra with antipode s over the field k . By Lemma 2 it follows that $R \in s(A) \otimes s(A)$. As a consequence of Proposition 1 we now have:

THEOREM 1. *Suppose that (A, R) is a quasitriangular Hopf algebra with antipode s over the field k . Write $R = \sum R^{(1)} \otimes R^{(2)}$ and let $u = \sum s(R^{(2)}) R^{(1)}$. Then u is invertible, and $s^2(a) = uau^{-1}$ for all $a \in A$.*

It may be of interest to note that (QT.5) can be expressed in terms of A -module actions. Define A -module actions on A by

$$a \rightarrow b = ab \quad \text{and} \quad b \leftarrow a = ba$$

for $a, b \in A$, on A^* by

$$a \rightarrow p(b) = p(b \leftarrow a) \quad \text{and} \quad p \leftarrow a(b) = p(a \rightarrow b)$$

for $a, b \in A$ and $p \in A^*$, and on $\text{Hom}_k(A^*, A)$ by

$$a \rightarrow f(p) = \sum a_{(1)} \rightarrow (f(p \leftarrow a_{(2)}))$$

and

$$f \leftarrow a(p) = \sum (f(a_{(1)} \rightarrow p)) \leftarrow a_{(2)}$$

for $a \in A$, $f \in \text{Hom}_k(A^*, A)$, and $p \in A^*$. Notice that $a \rightarrow (f \leftarrow b) = (a \rightarrow f) \leftarrow b$ for all $a, b \in A$ and $f \in \text{Hom}_k(A^*, A)$. Therefore

$$a \rightarrow f = \sum a_{(1)} \rightarrow f \leftarrow s(a_{(2)})$$

defines a left A -module action on $\text{Hom}_k(A^*, A)$. It is easy to see that the A -module actions \rightarrow and \leftarrow on $\text{Hom}_k(A^*, A)$ are related by

$$a \rightarrow f = \sum (a_{(1)} \rightarrow f) \leftarrow a_{(2)} \tag{7}$$

for all $a \in A$ and $f \in \text{Hom}_k(A^*, A)$. The proof of the following proposition is left to the reader.

PROPOSITION 2. *Suppose that A is a Hopf algebra over the field k and $R = \sum R^{(1)} \otimes R^{(2)} \in A \otimes A$. Define $f \in \text{Hom}(A^*, A)$ by $f(p) = \sum p(R^{(1)}) R^{(2)}$ for $p \in A^*$. Give $\text{Hom}_k(A^*, A)$ the A -module structures described above. Then the following are equivalent:*

- (a) $\sum a_{(2)} R^{(1)} \otimes a_{(1)} R^{(2)} = \sum R^{(1)} a_{(1)} \otimes R^{(2)} a_{(2)}$ for all $a \in A$,
- (b) $a \rightarrow f = f \leftarrow a$ for all $a \in A$,
- (c) $a \rightarrow f = \varepsilon(a) f$ for all $a \in A$.

3. s^4 WHEN (A, R) IS FINITE-DIMENSIONAL

Suppose that (A, R) is a finite-dimensional quasitriangular Hopf algebra with antipode s over the field k . Write $R = \sum R^{(1)} \otimes R^{(2)}$, let $u = \sum s(R^{(2)}) R^{(1)}$, and $v = s(u)^{-1}$. In this section we show that $h = vu$ is a grouplike element and $s^4(a) = hah^{-1}$ for all $a \in A$. Generally s^4 need not be inner in the finite-dimensional case [5]. The main results of this section are implications of equation (QT.5) specialized to $a = A$, where A is a non-zero left integral for A .

Suppose that A is any finite-dimensional Hopf algebra with antipode s over the field k . An element $A \in A$ is said to be a left (respectively right) integral if $aA = \varepsilon(a)A$ (respectively $AA = \varepsilon(a)A$) for all $a \in A$. Thus $\lambda \in A^*$ is a right (respectively left) integral if $\lambda p = p(1)\lambda$ (respectively $p\lambda = p(1)\lambda$) for all $p \in A^*$. The space of left (or right) integrals of A is a one-dimensional ideal. If $A \in A$ is a non-zero left (or right) integral, then A is a free A^* -module with basis A under the left action defined by $p \rightarrow a = \sum a_{(1)} p(a_{(2)})$ and also under the right action defined by $a \leftarrow p = \sum p(a_{(1)}) a_{(2)}$ for $p \in A^*$ and $a \in A$. Thus if $\lambda \in A^*$ is a non-zero right (or left) integral, then A^* is a free A -module with basis λ under both A -module actions on A^* defined in the last section. We direct the reader to [6, Chap. 5] for a basic treatment of integrals with proofs.

A non-zero $a \in A$ is called a grouplike element if $A(a) = a \otimes a$. If a is a grouplike element, then $\varepsilon(a) = 1$ and $s(a)$ is a grouplike element which is a multiplicative inverse of a . It is easy to see that the set of grouplike elements $G(A)$ of A is a group under multiplication, and that $G(A^*) = \text{Alg}_k(A, k)$.

Let $A \in A$ be a non-zero left integral. Since A generates a one-dimensional ideal of A , there is a unique algebra homomorphism $\alpha \in \text{Alg}_k(A, k) = G(A^*)$ such that $AA = \alpha(a)A$ for all $a \in A$. Now let $\lambda \in A^*$ be a non-zero right integral. Then there is a unique algebra

homomorphism $g \in \text{Alg}_k(A^*, k) = G(A^{**})$ such that $p\lambda = g(p)\lambda$ for all $p \in A^*$. We shall call $g \in A^{**} = A$ the distinguished grouplike element of A and α the distinguished grouplike element of A^* . By [3, Proposition 6]

$$s^4(a) = g(\alpha \rightarrow a \leftarrow \alpha^{-1}) g^{-1} \quad \text{for all } a \in A. \quad (8)$$

By [4, Theorem 2(c)]

$$\sum A_{(2)} \otimes A_{(1)} = \sum A_{(1)} \otimes s^2(A_{(2)}) g. \quad (9)$$

Let $\eta \in G(A^*) = \text{Alg}_k(A, k)$. We say that $a \in A$ is a left (respectively right) η -integral if $ba = \eta(b)a$ (respectively $ab = \eta(b)a$) for all $b \in A$.

LEMMA 3. *Suppose that A is a finite-dimensional Hopf algebra with antipode s over the field k , and let $\eta \in G(A^*)$. Then*

(a) *if $a \in A$ is a left η -integral, then $\sum a_{(1)} \otimes ba_{(2)} = \sum s(\eta \rightarrow b) a_{(1)} \otimes a_{(2)}$ for all $b \in A$,*

(b) *if $a \in A$ is a right η -integral, then $\sum a_{(1)} b \otimes a_{(2)} = \sum a_{(1)} \otimes a_{(2)} s(b \leftarrow \eta)$ for all $b \in A$.*

Proof. Let $a, b \in A$. The lemma follows directly from the equations $\sum a_{(1)} \otimes ba_{(2)} = \sum s(b_{(1)})(b_{(2)}a)_{(1)} \otimes (b_{(2)}a)_{(2)}$ and $\sum a_{(1)} b \otimes a_{(2)} = \sum (ab_{(1)})_{(1)} \otimes (ab_{(1)})_{(2)} s(b_{(2)})$. The derivation of the first equation is

$$\begin{aligned} \sum a_{(1)} \otimes ba_{(2)} &= \sum (\varepsilon(b_{(1)}) 1) a_{(1)} \otimes b_{(2)} a_{(2)} \\ &= \sum s(b_{(1)}) b_{(2)} a_{(1)} \otimes b_{(3)} a_{(2)} \\ &= \sum s(b_{(1)})(b_{(2)}a)_{(1)} \otimes (b_{(2)}a)_{(2)} \end{aligned}$$

and the similar derivation of the second is left to the reader. This concludes our proof.

Let (A, R) be a finite-dimensional quasitriangular Hopf algebra with antipode s over the field k . For $\eta \in G(A^*)$ define $g_\eta = \sum R^{(1)} \eta(R^{(2)})$, where $R = \sum R^{(1)} \otimes R^{(2)}$. There is an interesting relationship between the groups $G(A^*)$ and $G(A)$.

PROPOSITION 3. *Suppose that (A, R) is a finite-dimensional quasitriangular Hopf algebra with antipode s over the field k , and let $\eta \in G(A^*)$. Then*

(a) $g_\eta \in G(A)$,

(b) *the map $G(A^*) \rightarrow G(A)$ given by $\eta \mapsto g_\eta$ is a group anti-homomorphism,*

- (c) $(a \leftarrow \eta) g_\eta = g_\eta (\eta \rightarrow a)$ for all $a \in A$, and
- (d) η is central if and only if g_η is central.

Proof. We first show part (a). By (QT.1) it follows that $\Delta g_\eta = \sum \Delta R^{(1)} \eta(R^{(2)}) = \sum R^{(1)} \otimes r^{(1)} \eta(R^{(2)}) r^{(2)} = \sum R^{(1)} \otimes r^{(1)} \eta(R^{(2)}) \eta(r^{(2)}) = g_\eta \otimes g_\eta$ since η is an algebra homomorphism. By (QT.2) we have that $\varepsilon(g_\eta) = \eta(\sum \varepsilon(R^{(1)}) R^{(2)}) = \eta(1) = 1$. Therefore $g_\eta \in G(A)$, and part (a) is established.

We next show part (b). For $\eta, \rho \in G(A^*)$ we see by (QT.3) that $g_{\eta\rho} = \sum R^{(1)} \eta\rho(R^{(2)}) = \sum R^{(1)} \eta(R^{(2)}_{(1)}) \rho(R^{(2)}_{(2)}) = \sum R^{(1)} r^{(1)} \eta(r^{(2)}) \rho(R^{(2)}) = g_\rho g_\eta$.

To prove part (c), recall that $\sum a_{(2)} R^{(1)} \otimes a_{(1)} R^{(2)} = \sum R^{(1)} a_{(1)} \otimes R^{(2)} a_{(2)}$ for all $a \in A$ by (QT.5). Applying $I \otimes \eta$ to both sides of this equation gives part (c).

We finally prove part (d). First observe that $p(a \leftarrow q) = qp(a)$ and $p(q \rightarrow a) = pq(a)$ for all $p, q \in A^*$ and $a \in A$. Therefore $q \in A^*$ is central if and only if $a \leftarrow q = q \rightarrow a$ for all $a \in A$. Since η is invertible, it follows that $A \leftarrow \eta = A = \eta \rightarrow A$. Thus part (d) follows by part (c). The proof of the proposition is complete.

Part (c) of Proposition 3 has an interesting interpretation. Let $\eta \in G(A^*)$. Then $T_\eta: A^* \rightarrow A^*$ defined by $T_\eta(p) = \eta p \eta^{-1}$ for $p \in A^*$ is an automorphism of Hopf algebras. Observe that $T_\eta = t_\eta^*$ where $t_\eta: A \rightarrow A$ is defined by $t_\eta(a) = \eta^{-1} \rightarrow a \leftarrow \eta$ for $a \in A$. By part (c) of Proposition 3 we have that $t_\eta(a) = g_\eta a g_\eta^{-1}$ for $a \in A$. In particular t_η is itself inner.

If (A, R) is a finite-dimensional quasitriangular Hopf algebra with antipode s over the field k , then by the comments made in the preceding paragraph and Eq. (8) it follows that $s^4(a) = h a h^{-1}$ for all $a \in A$, where $h = g g_{x^{-1}}$. The grouplike element h can be computed directly from R .

THEOREM 2.¹ *Suppose that (A, R) is a finite-dimensional quasitriangular Hopf algebra over the field k . Write $R = \sum R^{(1)} \otimes R^{(2)}$, let $u = \sum s(R^{(2)}) R^{(1)}$, and $v = s(u)^{-1}$. Suppose that g is the distinguished grouplike element of A , that α is the distinguished grouplike element of A^* , and let $h = g g_{x^{-1}}$. Then*

- (a) $g = v(\sum s(R^{(2)} \leftarrow \alpha) R^{(1)})$,
- (b) $g = v u g_\alpha$, and
- (c) $h = v u$. Thus vu is a grouplike element, and $s^4(a) = (vu) a (vu)^{-1}$ for all $a \in A$.

¹ Part (c) and Corollary 2 below are derived in Section 3 of [DRIN]. Part (b) is Proposition 6.1 of [DRIN] for A^{op} , and is proved by different methods there. The author was made aware of [DRIN] after this paper was accepted.

Proof. We first show part (a). Let $\lambda \in A$ be a non-zero left integral. Then λ is a left ε -integral and a right α -integral. Since $\sum \lambda_{(2)} R^{(1)} \otimes \lambda_{(1)} R^{(2)} = \sum R^{(1)} \lambda_{(1)} \otimes R^{(2)} \lambda_{(2)}$ by (QT.5), it follows by Lemma 3 that

$$\sum \lambda_{(2)} s(R^{(2)} \leftarrow \alpha) R^{(1)} \otimes \lambda_{(1)} = \sum R^{(1)} s(R^{(2)}) \lambda_{(1)} \otimes \lambda_{(2)}.$$

By Eq. (9) we have that $\sum \lambda_{(1)} \otimes \lambda_{(2)} = \sum s^2(\lambda_{(2)}) g \otimes \lambda_{(1)}$. Therefore

$$\sum \lambda_{(2)} s(R^{(2)} \leftarrow \alpha) R^{(1)} \otimes \lambda_{(1)} = \sum R^{(1)} s(R^{(2)}) s^2(\lambda_{(2)}) g \otimes \lambda_{(1)}.$$

Since (A, \leftarrow) is a free right A^* -module with basis λ , there is some $p \in A^*$ such that $\lambda \leftarrow p = 1$. Thus applying $I \otimes p$ to both sides of the equation above we have that $\sum s(R^{(2)} \leftarrow \alpha) R^{(1)} = \sum R^{(1)} s(R^{(2)}) g$. Now $\sum R^{(1)} s(R^{(2)}) = \sum s(R^{(1)}) s^2(R^{(2)}) = s(u) = v^{-1}$ by Lemma 2. Therefore $\sum s(R^{(2)} \leftarrow \alpha) R^{(1)} = v^{-1}g$, and part (a) follows.

To establish part (b), we show that $\sum s(R^{(2)} \leftarrow \alpha) R^{(1)} = ug_\alpha$. Now $\sum R^{(1)} \otimes \lambda R^{(2)} = \sum R^{(1)} r^{(1)} \otimes r^{(2)} \otimes R^{(2)}$ by (QT.3). Therefore $\sum R^{(1)} \otimes R^{(2)} \leftarrow \alpha = \sum R^{(1)} r^{(1)} \alpha(r^{(2)}) \otimes R^{(2)} = \sum R^{(1)} g_\alpha \otimes R^{(2)}$. Hence $\sum s(R^{(2)} \leftarrow \alpha) R^{(1)} = \sum s(R^{(2)}) R^{(1)} g_\alpha = ug_\alpha$, and part (b) is established.

To show part (c), we first note that $vu = g(g_\alpha)^{-1} = gg_{\alpha^{-1}}$ by part (b) of Proposition 3 and part (b) above. Therefore $h = vu$ and is a grouplike element. By Eq. (5) it follows that $s^4(a) = hah^{-1}$ for all $a \in A$. Thus part (c) follows, and the proof of the theorem is complete.

A finite-dimensional Hopf algebra is said to be unimodular if $\alpha = \varepsilon$, or equivalently left and right integrals for A are the same. Thus by part (a) of Theorem 2 we have:

COROLLARY 1. *Suppose that (A, R) is a finite-dimensional quasi-triangular Hopf algebra over the field k . If A is unimodular, then $g = vu$, where g is the distinguished grouplike element of A .*

The Quantum Casimir of a quasitriangular Hopf algebra (A, R) over k is defined [2, p. 66] to be the element $c = us(u)$, where u is described in the preceding theorem. As a corollary to part (c) of Theorem 2 we have:

COROLLARY 2. *Suppose that (A, R) is a finite-dimensional quasi-triangular Hopf algebra over the field k . Write $R = \sum R^{(1)} \otimes R^{(2)}$ and let $u = \sum s(R^{(2)}) R^{(1)}$. Then the Quantum Casimir $c = u^2 l$ for some grouplike element $l \in G(A)$.*

If (A, R) is a finite-dimensional quasitriangular Hopf algebra over the field k , then the product vug_α does not depend on R by part (b) of Theorem 2. The element u may very well depend on R , as an easy example shows.

Suppose that $n > 1$ and that k has a primitive n^{th} root of unity ω . Let $A = k[G]$ be the group algebra of the cyclic group $G = \langle a \rangle$ of order n over k . Then $(A, 1 \otimes 1)$ is quasitriangular since A is cocommutative. For this particular structure $u = 1$.

Now set

$$R = \frac{1}{n} \left(\sum_{0 \leq l, m < n} \omega^{-lm} a^l \otimes a^m \right).$$

Then (A, R) is quasitriangular. To see this, let $\eta \in G(A^*)$ be the algebra homomorphism determined by $\eta(a) = \omega$. Since distinct grouplike elements are linearly independent [6, Proposition 3.2.1.b)], it easily follows that the cyclic group $G' = \langle \eta \rangle$ has order n , and thus $A^* = k[G']$ is the group algebra of G' over k . Now $f: A^* \rightarrow A$ defined by $f(\eta^l) = a^l$ is a bialgebra isomorphism. Using the equation $\sum_{l=0}^{n-1} \omega^{ul} = n\delta_{0,u}$ for $u = 0, \dots, n-1$ it is easy to see that $f = F(R)$, where $F: A \otimes A \rightarrow \text{Hom}_k(A^*, A)$ is the identification defined in Section 2. Therefore (QT.1)–(QT.4) are satisfied. Since A is commutative and cocommutative (QT.5) is satisfied also. Thus (A, R) is quasitriangular.

For this particular R it is easy to see that

$$u = \frac{1}{n} \left(\sum_{0 \leq m, l < n} \omega^{l(m-l)} a^m \right).$$

If $n = 2$ then $u = a$. Suppose that $n = 3$. Then

$$u = \frac{1}{3} ((1 + 2\omega^2) 1 + (2 + \omega) a + (2 + \omega) a^2).$$

In this case note that u and the Quantum Casimir

$$c = \frac{1 - \omega}{3} (\omega 1 + a + a^2)$$

are not grouplike elements.

We conclude this paper by finding a necessary and sufficient condition for s^2 to be inner, where s is the antipode of a finite-dimensional Hopf algebra over k , in the spirit of Proposition 1. The proposition below is motivated by the proof of part (a) of Theorem 2.

PROPOSITION 4. *Let A be a finite-dimensional Hopf algebra with antipode s over the field k . Suppose that $\lambda \in A$ is a non-zero left integral. Then the following are equivalent:*

- (a) s^2 is inner,
- (b) there are $R, U \in A \otimes A$ such that $\sum \lambda_{(2)} \otimes \lambda_{(1)} = \sum R(\lambda_{(1)} \otimes \lambda_{(2)}) U$.

Proof. We first show that part (a) implies part (b). Suppose that $u \in A$ is invertible and that $s^2(a) = uau^{-1}$ for all $a \in A$. By Eq. (9) we have $\sum A_{(2)} \otimes A_{(1)} = \sum A_{(1)} \otimes s^2(A_{(2)}) g$, where g is the distinguished grouplike element of A . Let $R = 1 \otimes u$ and $U = 1 \otimes u^{-1}g$.

We show part (b) implies part (a). Assume that $R, U \in A \otimes A$ satisfy $\sum A_{(2)} \otimes A_{(1)} = \sum R(A_{(1)} \otimes A_{(2)}) U$. Write $R = \sum R^{(1)} \otimes R^{(2)}$ and $U = \sum U^{(1)} \otimes U^{(2)}$. Then by Lemma 3 we have that

$$\begin{aligned} \sum R(A_{(1)} \otimes R_{(2)}) U &= \sum R^{(1)} A_{(1)} U^{(1)} \otimes R^{(2)} A_{(2)} U^{(2)} \\ &= \sum A_{(1)} \otimes R^{(2)} s^{-1}(R^{(1)}) A_{(2)} s(U^{(1)} \leftarrow \alpha) U^{(2)}, \end{aligned}$$

where α is the distinguished grouplike element of A^* . Hence $\sum A_{(2)} \otimes A_{(1)} = \sum A_{(1)} \otimes u A_{(2)} v$ for some $u, v \in A$. But by Eq. (9) again, we have that $\sum A_{(1)} \otimes s^2(A_{(2)}) g = \sum A_{(1)} \otimes u A_{(2)} v$. This last equation is equivalent to $s^2(A \leftarrow p) g = u(A \leftarrow p) v$ for all $p \in A^*$. Since (A, \leftarrow) is a free right A^* -module with basis A , this last equation implies that $s^2(a) g = uav$ for all $a \in A$. Therefore u is invertible and has inverse vg^{-1} , and hence $s^2(a) = uau^{-1}$ for all $a \in A$. This concludes the proof of the proposition.

REFERENCES

1. V. G. DRINFEL'D, Quantum groups, in "Proceedings of the International Congress of Mathematicians, Berkeley, CA, 1987," pp. 798–820.
2. V. G. DRINFEL'D, On almost cocommutative Hopf algebra, *Leningrad Math. J.* **1** (1990), 321–342.
3. SHAHN MAJID, Quasitriangular Hopf algebras and Yang–Baxter equations, *Internat. J. Modern Phys. A* **5** (1990), 1–91.
4. D. E. RADFORD, The order of the antipode of a finite-dimensional Hopf algebra is finite, *Amer. J. Math.* **98** (1976), 333–355.
5. D. E. RADFORD, The trace function and finite-dimensional Hopf algebras, preprint, 1990.
6. D. E. RADFORD, On the coradical of a finite-dimensional Hopf algebra, *Proc. Amer. Math. Soc.* **53** (1975), 9–15.
7. M. E. SWEEDLER, "Hopf Algebras," Benjamin, New York, 1969.